

学校编码: 10384
学号: 21620100153876

分类号____密级____
UDC____

厦门大学
博士学位论文

菲和芘的心脏发育毒性及其机制的研究

Effects of phenanthrene and pyrene on cardiac
development of embryonic zebrafish (*Danio rerio*): the
mechanism involved

章幼玉

指导教师姓名: 王重刚 教授

专业名称: 动物学

论文提交日期: 2013 年 5 月

论文答辩时间: 2013 年 6 月

学位授予日期:

答辩委员会主席: _____
评阅人: _____

2013 年 5 月

菲和芘的心脏发育毒性及其机制的研究

章幼玉

指导教师: 王重刚 教授

厦门大学

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

- () 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。
- () 2. 不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

目 录

目 录	I
CONTENTS	V
摘要	VIII
ABSTRACT	X
第一章 前言	1
1.1 多环芳烃污染及研究现状	1
1.1.1 多环芳烃的理化特性	1
1.1.2 多环芳烃的污染现状	2
1.2 多环芳烃的毒性效应及机制研究	4
1.2.1 致癌性、致畸性和致突变性	4
1.2.2 胚胎发育毒性	5
1.2.3 其它毒性	6
1.3 鱼类心脏发育的过程和调控机制	7
1.3.1 斑马鱼心脏发育过程	7
1.3.2 鱼类心脏发育的调控机制	8
1.3.3 鱼类心脏功能执行相关的调控机制	12
1.4 多环芳烃对鱼类心脏发育的影响及其研究进展	13
1.5 本研究的内容、目的和意义	15
第二章 材料与amp;方法	16
2.1 实验材料	16
2.2 主要仪器	16
2.3 主要药品与试剂	17
2.4 合成及测序	17
2.5 主要试剂配制	17

2.5.1 斑马鱼胚胎收集培养及显微注射所需主要溶液	17
2.5.2 DNA 片段胶回收及克隆所需主要溶液	18
2.5.3 大肠杆菌感受态制备相关溶液	18
2.5.4 大肠杆菌培养培养基	18
2.5.5 质粒提取所需主要溶液	18
2.5.6 Western blot 分析所需主要溶液	18
2.5.7 石蜡切片及 HE 染色所需主要溶液	19
2.5.8 Masson 三色染色法所需主要溶液	19
2.5.9 原位杂交所需主要溶液	19
2.6 主要实验方法	20
2.6.1 毒物暴露实验	20
2.6.2 斑马鱼心脏发育过程中相关形态学和功能等指标的毒物学效应观察	20
2.6.3 石蜡包埋及染色	21
2.6.4 基因芯片筛选差异基因	21
2.6.5 组织总 RNA 提取	22
2.6.6 1st cDNA 合成	23
2.6.7 基因克隆	23
2.6.8 实时荧光定量 PCR (Realtime-PCR, RT-PCR) 分析	25
2.6.9 原位杂交	27
2.6.10 Western blot	30
2.6.11 免疫组化	31
2.6.12 明胶酶谱分析	31
2.6.13 H9C2 细胞培养及其相关实验	31
2.7 数据处理	32
第三章 结果分析	33
3.1 菲、芘 暴露对斑马鱼胚胎心脏发育的影响	33
3.1.1 Phe、Pyr 暴露对心脏形态形成的影响	33
3.1.2 Phe、Pyr 暴露对心脏功能的影响	38

3.2 菲暴露对斑马鱼胚胎心脏发育影响的机制	42
3.2.1 Phe 暴露对心血管发育或疾病相关调控基因的影响	42
3.2.2 Phe 影响斑马鱼胚胎心脏形态形成的分子机制	44
3.2.2.1 Phe 暴露对斑马鱼胚胎心脏发育及功能相关的基因表达的影响	44
3.2.2.2 Phe 暴露对 MMP-9 表达及活性的影响	49
3.2.2.3 MMP-9 inhibitor 缓解了 Phe 诱发的心脏畸形	51
3.2.2.4 Phe 暴露对胶原合成的影响	53
3.2.3 Phe 影响斑马鱼心脏功能的分子机制	56
3.2.3.1 Phe 暴露对大鼠心肌细胞钙离子平衡的影响	56
3.2.3.2 Phe 暴露对大鼠心肌细胞钙离子通道相关蛋白的影响	57
3.2.3.3 Phe 暴露对斑马鱼胚胎钙泵蛋白的影响	60
3.2.3.4 Phe 暴露对 TBX5 表达的影响	63
3.3 芘暴露对斑马鱼胚胎心脏发育影响的机制	65
3.3.1 Pyr 暴露对 AhR 和 Cyp1a 基因表达的影响	65
3.3.2 Pyr 暴露对 MMP-9 表达和活性的影响	67
3.3.3 Pyr 暴露对心脏发育相关基因表达的影响	69
3.3.4 Pyr 暴露对心脏功能执行相关基因表达的影响	71
第四章 讨论	72
4.1 菲、芘暴露引发心脏发育畸形和心脏功能障碍	72
4.2 菲暴露干扰心脏发育和心脏功能执行相关的分子	73
4.2.1 Phe 暴露引发心脏发育缺陷和心脏功能障碍的分子机制	74
4.2.1.1 Phe 暴露引发心脏发育缺陷的分子机制	74
4.2.1.2 Phe 暴露引发心率不齐的分子机制	76
4.2.2 Pyr 暴露引发心脏发育缺陷的分子机制	78
第五章 结论	81
5.1 本研究所取得的主要结果	81
5.2 本研究的创新点	81
5.3 展望	82

参考文献	83
缩略词对照表	98
攻读博士阶段发表的论文和参加的课题	100
致 谢	101

厦门大学博硕士论文摘要库

CONTENTS

Abstract (In Chinese)	VIII
Abstract (In English)	X
Chapter 1 Introduction	1
1.1 The study of PAHs polltion	1
1.1.1 The characters of PAHs	1
1.1.2 The pollution study of PAHs	2
1.2 The toxicities of PAHs	4
1.2.1 Carcinogenicity	4
1.2.2 Development toxicity	5
1.2.3 others	6
1.3 The process of heart development in fish and its mechanism	7
1.3.1 The process of heart development in zebrafish	7
1.3.2 The mechanisms of heart development in fish	8
1.3.3 The mechanisms of functional heart regulation in fish	12
1.4 The effects of PAHs on heart development	13
1.5 Research objectives and significance	15
Chapter 2 Materials and methods	16
2.1 Materials	16
2.2 Apparatus	26
2.3 Main regents	17
2.4 Sequencing	17
2.5 Solutions	17
2.6 Mthods	20
2.6.1 Exposure experiment	20
2.6.2 Assessment of cardiac toxicity	20

2.6.3 Paraffin-embedded, section and staining	21
2.6.4 Microarray	21
2.6.5 Total RNA extraction	22
2.6.6 cDNA synthesization	23
2.6.7 Gene cloning	23
2.6.8 Realtime-PCR assay	25
2.6.9 <i>In situ</i> hybridization	27
2.6.10 Western blot analysis	30
2.6.11 Immunohistochemistry	31
2.6.12 Gelatin zymography	31
2.6.13 Cell culture and the measurement of calcium transients	31
2.7 Data analysis	32
Chapter 3 Results	33
3.1 Effects of Phe and Pyr on cardiac development of zebrafish embryos ..	33
3.1.1 Effects of Phe and Pyr on heart morphology	33
3.1.2 Effects of Phe and Pyr on cardiac function	38
3.2 Effects of Phe on heart development and the machenism involved	42
3.2.1 Effects of Phe on the expression of cardiac development correlated genes	42
3.2.2 The machanism of cardiac toxicity of Phe	44
3.2.2.1 Effects of Phe on the expression of cardiac development or function correlated genes	44
3.2.2.2 Effects of Phe on the expression of MMP-9	49
3.2.2.3 MMP-9 inhibitor attenuated the cardiac defects caused by Phe ..	51
3.2.2.4 Effects of Phe on collagen synthesis in the ventricle	53
3.2.3 The mechanism underlying Phe induced cardiac dysfunction	56
3.2.3.1 Effects of Phe on Ca ²⁺ handling in cardiac myoblasts	56
3.2.3.2 Effects of Phe on Ca ²⁺ -cycling channels	57

3.2.3.3 Effects of Phe on the expression of SERCA2a in zebrafish larvae	60
3.2.3.4 Effects of Phe on the expression of TBX5	63
3.3 Effects of Pyr on heart development and the mechanism involved.....	65
3.3.1 Effects of Pyr on the expression of AhRs and Cyp1a	65
3.3.2 Effects of Pyr on the expression of MMP-9	67
3.3.3 Effects of Pyr on the expression of cardiac development relevant genes	69
3.3.4 Effects of Pyr on the expression of cardiac function correlated genes ..	71
Chapter 4 Discussion	72
4.1 Embryonic exposure of Phe or Pyr induced cardiac defects and dysfunction.....	72
4.2 Exposure of Phe disturbed the expression of cardiac development and function relevant genes	73
4.2.1 The mechanism of cardiac toxicity of Phe	74
4.2.1.1 The mechanism underlying heart defects caused by Phe	74
4.2.1.2 The mechanism of arrhythmia caused by Phe.....	76
4.2.2 The mechanism of cardiac toxicity of Pyr.....	78
Chapter 5 Conclusions	81
References	83
Abbreviation	98
Publications and Projects	100
Acknowledgement.....	101

摘要

多环芳烃 (Polycyclic aromatic hydrocarbons, PAHs) 是一类广泛分布于环境中的持久性有机污染物 (persistent organic pollutants, POPs)。众多研究表明, PAHs 不仅具有较强的致癌、致畸和致突变性, 还具有胚胎发育毒性, 尤其是心脏发育毒性。但到目前为止, PAHs 心脏毒性的研究多集中于高浓度的急性暴露, 而对环境浓度的 PAHs 暴露是否会同样的引发心脏发育毒性, 研究还甚少; 且关于低环数的 PAHs 的心脏发育毒性机制研究尚不清楚。本研究分别选取 3 环的菲 (phenanthrene, Phe) 和 4 环的芘 (pyrene, Pyr), 在环境水平 (0, 0.05, 5, 50 nM) 暴露斑马鱼 (*Danio rerio*) 胚胎, 应用生态毒理学和分子生物学技术检测在胚胎发育过程中其对心脏发育的毒性效应, 旨在探讨: (1) 环境浓度的 Phe、Pyr 暴露是否同样会引发心脏发育缺陷; (2) 比较 Phe、Pyr 暴露对鱼类心脏发育毒性效应的差异; (3) 比较分析其可能的作用机制。结果显示:

1) 斑马鱼受精卵在环境水平的 Phe 暴露下, 在胚胎发育过程中, 各处理组心脏均产生一定的畸形, 且总的心脏畸形率呈显著性上升, 主要畸形特征表现为: 心包囊水肿、心房与心室重叠减少、心室扩大等。通过 HE 染色等组织学方法分析显示, Phe 暴露能引起斑马鱼胚胎心室腔变大, 心室壁变薄, 心室心肌细胞层胶原含量增加等畸形。同时, 与心脏形态形成相关的 MMP-9 基因, 不论是转录水平、蛋白水平还是蛋白活性均呈显著性上升, 且通过使用 MMP-9 抑制剂特异性的抑制 MMP-9 活性, Phe 暴露引发的心脏发育缺陷在很大程度上得到了缓解。另外, 心室细胞层中, 在心肌纤维化过程中与胶原合成密切相关的调控因子 TGF- β 的表达也呈显著性上升。这些结果表明, Phe 暴露可能通过诱导 MMP-9 表达、增加 MMP-9 活性, 促进心肌细胞间质 ECM 降解、导致腔室壁变薄; Phe 暴露同时诱发 TGF- β 1 调节的胶原合成的增加, 最终导致心脏畸形。

2) 环境浓度的 Phe 暴露不仅对心脏形态发育产生不良影响, 同时也导致心脏机能障碍。Phe 暴露 72h, 各处理组心跳速度呈显著性上升, 且表现出严重的不规律性, 舒张/收缩末期容积 (EDV、ESV) 上升, 每搏输出量 (SV) 下降, 心输出量 (CO) 减少等。通过体外暴露大鼠心肌细胞研究发现, Phe 暴露导致

细胞胞浆内 Ca^{2+} 滞留严重, Ca^{2+} 浓度呈显著性上升, 而肌浆网内储存的 Ca^{2+} 则明显减少; 肌浆网钙泵 ATP 酶 SERCA2a 基因转录水平、蛋白表达及其活性均显著性下降; 另外, SERCA2a 的一个重要调节因子 Tbx5 的表达也显著性下降。这些结果表明, Phe 可能通过抑制 Tbx5 表达, 使得 SERCA2a 表达降低, 心肌细胞肌浆网对胞浆中 Ca^{2+} 摄取减少, 胞浆中 Ca^{2+} 滞留而扰乱了心肌细胞内钙离子平衡, 最终导致收缩、舒张紊乱并引发心率不齐。

3) 环境浓度的 Pyr 暴露引发与 Phe 类似的心脏发育毒性, 如: 心包囊水肿, 心脏非正常环化, 心室壁变薄, 心室细胞层胶原含量增加, 心跳加快等。但是 Pyr 暴露 72h, EDV、ESV 均表现出不同程度的下降趋势, 且都在 50 nM 呈显著性下降, SV 在 50 nM 处理组显著性减小, CO 则没有发生明显的变化。然而, 在作用机制上 Pyr 和 Phe 却不尽相同, Pyr 暴露并不会引起斑马鱼胚胎 MMP-9 基因表达的变化, Bmp2b, Nkx2.5 基因的表达却都呈显著性下降。因此, Pyr 暴露可能是通过影响 Bmp2b 的表达调节 Nkx2.5, 从而影响心脏心肌细胞的分化, 进而导致心脏发育缺陷。

综上所述, 本研究表明环境浓度 Phe 和 Pyr 暴露均能引发类似的心脏发育毒性, 但在作用机制上两者却不尽相同。Phe 主要是通过 MMP-9 介导的相关途径诱发心脏发育缺陷和通过扰乱心肌细胞 Ca^{2+} 平衡引发心脏功能障碍, 而 Pyr 可能是通过 Bmp2b-Nkx2.5 途径而发生作用的。

关键词: 多环芳烃 斑马鱼 胚胎发育 心脏 钙离子 机制

ABSTRACT

Polycyclic aromatic hydrocarbons (PAHs) are a kind of the most abundant and ubiquitous persistent organic pollutants (POPs) in environment. It is widely accepted that PAHs not only have carcinogenic, teratogenic, and mutagenic toxicity, but also developmental toxicity, especially the cardiotoxic effects. However, most of the studies concerning PAHs focus on high dose exposure, and the effects of exposure embryos to environmentally relevant levels of PAHs are still limited. In the present study, zebrafish (*Danio rerio*) embryos were exposed to Phe (phenanthrene) and Pyr (pyrene) at doses of 0, 0.05, 0.5, 5 and 50 nM, and the aims of this thesis were: (1) whether exposure to environmentally relevant levels of Phe and Pyr could cause cardiac defects; (2) What are the differences of cardiac toxicity between Phe and Pyr, and (3) What are the mechanism(s) underlying. The results were as follows:

1) Exposure to environmentally relevant levels of Phe (0.05-50 nmol/L) induced a series of cardiac defects, which were characterized by abnormally looped and enlarged hearts, dilated and thinner ventricular walls, and increased interstitial fibrosis, and the frequency of these abnormalities increased in a dose-dependent manner. Meanwhile, the mRNA and protein expression levels of matrix metalloproteinase-9 (MMP-9), as well as the MMP-9 activity, were induced. Moreover, during co-treatment of the zebrafish embryos with MMP-9 inhibitor, the cardiac defects caused by Phe were attenuated. In addition, Phe exposure led to an up-regulation of transforming growth factor β (TGF- β), which plays a crucial role in mediating cardiac fibrosis. Taken together, our data indicated that the exposure to Phe of zebrafish embryos disrupted normal cardiac development, and that the cardiac defects induced by Phe were mediated by the MMP-9, while TGF- β was also involved in these cardiac defects.

2) Phe exposure also affected cardiac function of zebrafish larvae. By 72hpf (hours

post-fertilization), the heart rate was dramatically increased, which together with significant arrhythmia, both the EDV (End-diastolic volume) and ESV (End-systolic volume) of the ventricle were also increased, and the SV (stroke volume) and CO (cardiac output) were decreased. Meanwhile, a disordered calcium (Ca^{2+}) handling characterized by impaired sarcoplasmic reticulum (SR) Ca^{2+} uptake and obvious Ca^{2+} accumulation in cytoplasm was observed in rat embryonic cardiac myoblasts (H9C2) exposed to Phe. The mRNA level as well as protein expression of SERCA2a Ca^{2+} pump in zebrafish hearts or H9C2 cells were significantly decreased by Phe exposure. The activity of Ca^{2+} -ATPase in H9C2 cells was inhibited by Phe. Both the mRNA and protein level of TBX5, a direct regulator of SERCA2a, were significantly decreased by Phe exposure. These results suggested that the exposure to Phe could lead to irregular cardiac rhythm in zebrafish embryos via perturbing calcium handling pathway.

3) Pyr exposure also caused cardiac defects in zebrafish larvae. By 72 hpf, Pyr-treated embryos showed dose-dependent heart abnormalities, such as pericardial edema, cardiac looping defects and increased interstitial fibrosis. They also showed an abnormal contraction including increase of heart beat and decreased EDV, ESV and SV. However, the expression of MMP-9 and the activities were not obviously changed in response to Pyr exposure. The homeodomain transcription factor Nkx2.5 and the bone morphogenetic protein 2b (Bmp2b) were down-regulated. Taken together, these data indicated that embryonic exposure to low-level environmental Pyr disrupt normal cardiac development and alter expression of defective cardiac differentiation related genes.

In summary, the most important finding in our study was that embryonic exposure to both Phe and Pyr could cause some similar cardiac defects. However, the underlying mechanisms of cardiac toxicity were different: The cardiac defects induced by Phe were mediated by the MMP-9, and Phe also could lead to irregular cardiac rhythm via perturbing calcium handling pathway. And Pyr exposure might be through a Bmp2b-Nkx2.5 mediated pathway.

Keywords: PAHs, zebrafish, embryo development, heart, calcium, mechanism

厦门大学博硕士论文摘要库

第一章 前言

1.1 多环芳烃污染及研究现状

1.1.1 多环芳烃的理化性质

多环芳烃 (Polycyclic aromatic hydrocarbons, PAHs) 是一类由 2 个或 2 个以上苯环以线性排列、弯接或簇聚的方式连接而成的碳氢化合物，主要来源于煤、碳、石油、木材、有机高分子化合物等有机物的不完全燃烧或燃解过程，是重要的环境和食品污染物。迄今已发现有 200 多种 PAHs，图 1-1 中列出的为 16 种常见的 PAHs，其中有相当部分具有致癌性，如苯并(a)芘 [benzo(a)pyrene, B(a)P]，苯并(a)蒽 (benzanthracene, BaA) 等。PAHs 大部分是无色或淡黄色的结晶，个别具深色，熔点及沸点较高，蒸气压很小，水溶性较差，脂溶性较强，可在生物体内蓄积，能溶于丙酮、苯、二氯甲烷等有机溶剂。




图 1-1 16 种常见的 PAHs

Fig. 1-1 16 kinds of PAHs

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库